

Controlling High Blood Pressure CMS165v13

Last modified on 01/26/2026 2:22 pm EST

Measure Information

Description

Percentage of patients 18-85 years of age who had a diagnosis of essential hypertension starting before and continuing into, or starting during the first six months of the measurement period, and whose most recent blood pressure was adequately controlled (<140/90mmHg) during the measurement period.

Guidance

In reference to the numerator element, only blood pressure readings performed by a clinician or an automated blood pressure monitor or device are acceptable for numerator compliance with this measure. This includes blood pressures taken in person by a clinician and blood pressures measured remotely by electronic monitoring devices capable of transmitting the blood pressure data to the clinician. Blood pressure readings taken by an automated blood pressure monitor or device and conveyed by the patient to the clinician are also acceptable. It is the clinician's responsibility and discretion to confirm the an automated blood pressure monitor or device used to obtain the blood pressure is considered acceptable and reliable and whether the blood pressure reading is considered accurate before documenting it in the patient's medical record.

Do not include BP readings taken during an acute inpatient stay or an emergency department (ED) visit.

If no blood pressure is recorded during the measurement period, the patient's blood pressure is assumed "not controlled".

If there are multiple blood pressure readings on the same day, use the lowest systolic and the lowest diastolic reading as the most recent blood pressure reading. Ranges and thresholds do not meet criteria for this measure. A distinct numeric result for both the systolic and diastolic BP reading is required for numerator compliance.

This eCQM is a patient-based measure.

This version of the eCQM uses QDM version 5.6. Please refer to the eCQI resource center (<https://ecqi.healthit.gov/qdm>) for more information on the QDM.

Initial Population

Patients 18-85 years of age by the end of the measurement period who had a visit during the measurement period and diagnosis of essential hypertension starting before and continuing into, or starting during the first six months of the measurement period.

Denominator

Equals initial population

Denominator Exclusions

Exclude patients who are in hospice care for any part of the measurement period.

Patients with evidence of end stage renal disease (ESRD), dialysis or renal transplant before or during the measurement period. Also exclude patients with a diagnosis of pregnancy during the measurement period.

Exclude patients 66-80 by the end of the measurement period with an indication of frailty for any part of the measurement period who also meet any of the following advanced illness criteria:

- Advanced illness diagnosis during the measurement period or the year prior
- OR taking dementia medications during the measurement period or the year prior

Exclude patients 81 and older by the end of the measurement period with an indication of frailty for any part of the measurement period.

Exclude patients 66 and older by the end of the measurement period who are living long term in a nursing home any time on or before the end of the measurement period.

Exclude patients receiving palliative care for any part of the measurement period.

Denominator Exceptions

None

Numerator

Patients whose most recent blood pressure is adequately controlled (systolic blood pressure less than 140 mmHg and diastolic blood pressure less than 90 mmHg) during the measurement period.

Numerator Exclusions

None
